

ORAL PRESENTATION

Open Access

# Role of sarcoplasmic reticulum junctional proteins in skeletal muscle strength

Barbara Mosca<sup>1</sup>, Osvaldo Delbono<sup>2</sup>, Maria Laura Messi<sup>2</sup>, Leda Bergamelli<sup>1</sup>, Mirko Vukcevic<sup>3</sup>, Ruben Lopez<sup>1</sup>, Susan Treves<sup>1,3</sup>, Miyuki Nishi<sup>4</sup>, Hiroshi Takeshima<sup>4</sup>, Francesco Zorzato<sup>1,3\*</sup>

From 33rd Annual Meeting of the European Malignant Hyperthermia Group (EMHG)  
Würzburg, Germany. 15-17 May 2014

## Background

Skeletal muscle constitutes approximately 40% of body mass, and age-induced decrease of muscle strength impinge on daily activities and on normal social life in the elderly. Loss of muscle strength has been recognised as a debilitating and life threatening condition also in cachexia in cancer patients and in clinical conditions associated with prolonged bed rest. Skeletal muscle dihydropyridine receptors (Cav1.1) act as Ca<sup>2+</sup> channels and voltage sensors to initiate muscle contraction by activating ryanodine receptors, the Ca<sup>2+</sup> release channels of the sarcoplasmic reticulum. Cav1.1 activity is enhanced by a retrograde stimulatory signal delivered by the ryanodine receptor. JP45 is a membrane protein interacting with Cav1.1 and the sarcoplasmic reticulum Ca<sup>2+</sup> storage protein calsequestrin (CASQ1).

We hypothesized that JP45 and CASQ1 form a signaling pathway which modulates Cav1.1 channel activity.

## Materials and methods

We isolated flexor digitorum brevis (FDB) muscle fibres from JP45 and CASQ1 double knock-out mice (DKO) and tested whether there were differences in Ca<sup>2+</sup> homeostasis between the different mouse lines.

## Results

Our results show that Ca<sup>2+</sup> transients evoked by tetanic stimulation in DKO fibres, result from massive Ca<sup>2+</sup> influx due to enhanced Cav1.1 channel activity. This enhanced activity causes an increase of muscle strength both *in vitro* and *in vivo*.

## Conclusions

We conclude that skeletal muscle contraction is strengthened through the modulation of Cav1.1 channel activity by JP45 and CASQ1.

## Acknowledgements

This work was supported by funds from Swiss Muscle foundation, A.F.M., S.N. F and Department of Biomedicine University Hospital Basel. This study was also supported by Research Grant no. GGP08153 from the Italian Telethon ONLUS Foundation to F.P. and grants from the NIH/NIA (AG13934 and AG15820) to O.D.

## Authors' details

<sup>1</sup>Department of Life Sciences and Biotechnology, General Pathology section, University of Ferrara, Ferrara, 44100, Italy. <sup>2</sup>Department of Physiology and Pharmacology, Wake Forest University School of Medicine, Winston-Salem, N. C. 27157, USA. <sup>3</sup>Departments of Biomedicine and Anesthesiology, Basel University Hospital, Basel, 4031, Switzerland. <sup>4</sup>Department of Biological Chemistry, Graduate School of Pharmacological Sciences, Kyoto University, Tokyo, 113-0033, Japan.

Published: 18 August 2014

doi:10.1186/1471-2253-14-S1-A19

**Cite this article as:** Mosca et al.: Role of sarcoplasmic reticulum junctional proteins in skeletal muscle strength. *BMC Anesthesiology* 2014 **14**(Suppl 1):A19.

<sup>1</sup>Department of Life Sciences and Biotechnology, General Pathology section, University of Ferrara, Ferrara, 44100, Italy  
Full list of author information is available at the end of the article